

NOTE ON THE FLUX FOR TASEP WITH GENERAL DISORDER

ALLAN SLY

ABSTRACT. Extending results of Bahadoran and Bodineau we show that the flux rate of TASEP with independent and identically distributed disorder always has a plateau of densities around $\frac{1}{2}$.

The TASEP model (totally asymmetric simple exclusion process) on $\mathbb{Z}$ is a model of exclusion particles which always move to the right. Configurations in $\{0, 1\}^{\mathbb{Z}}$ represent the indicator of particles being present at each site on the line. In TASEP with rate r particles at site i jump to the right according to a rate r Poisson clock provided there is no particle present at site $i + 1$. A product measure with Bernoulli(ρ) is stationary for all ρ and the flux rate (the long term rate at which particles move across a bond) is given by

$$f(\rho) = r\rho(1 - \rho).$$

A natural problem is to consider the model where the jump rates of the bonds are chosen randomly from some law. In this model at bond i if a particle is present and $i + 1$ is unoccupied then the particle jumps from i to $i + 1$ according to a rate $\alpha(i)$ Poisson clock for independent and identically distributed variables $\alpha(i)$. This model was analysed in [1] in the case of a unit jump rate where an ϵ fraction of the sites have their jump rate replaced by some distribution $Q(\cdot)$ supported on $[r, 1]$. They show that the flux has a plateau around $\frac{1}{2}$, that is for some $0 < \rho_c < \frac{1}{2}$ it satisfies

$$f(\rho) = r/4$$

for all $\rho \in [\rho_c, 1 - \rho_c]$ provided ϵ is sufficiently small and Q satisfies condition **H** which says that as $u \searrow 0$ then

$$Q([r, r + u]) = O(u^\kappa),$$

for some $\kappa > 1$. They conjecture that this condition is not required and that the plateau should exist for any choice of Q and in particular the most natural choice of Q which is a point mass at r . The proof makes use of a sophisticated multi-scale analysis.

In this note we show that the most general version of the statement can be proved significantly more simply. Indeed if the jump rates α_i are independent and identically distributed and chosen from any positive non-constant distribution with infimum $r > 0$, i.e.

$$r = \inf\{s : \mathbb{P}[\alpha \geq s] < 1\}$$

then TASEP with site disorder α has a plateau.

Theorem 1. *With disorder α and $\mu = \frac{1}{r} - \mathbb{E}[\frac{1}{\alpha(i)}]$ for all $\rho \in [\frac{1}{2} - \frac{1}{4}\mu r, \frac{1}{2} + \frac{1}{4}\mu r]$,*

$$f(\rho) = r/4.$$

The simpler direction is the upper bound on the flux of $r/4$ which follows from the existence of wide bands of sites with disorder close to r as shown in Proposition 2.1 of [1].

Now consider the lower bound on the flux. The last passage percolation representation of TASEP is given as follows. For $\mathcal{W} = \{(i, j) \in \mathbb{Z}^2 : j \geq 0, i + j \geq 0\}$ let Y_{ij} be independent rate $\alpha(i)$ exponential random variables. A path γ from x to y in $\mathcal{W}$ is a sequence of vertices $x = (i_0, j_0), \dots, (i_k, j_k) = y$ in $\mathcal{W}$ such that

$$(i_\ell - i_{\ell-1}, j_\ell - j_{\ell-1}) \in \{(1, 0), (-1, 1)\}.$$

That is each step moves either east or northwest. For $(i, j) \in \mathcal{W}$ we write $\Gamma_{i,j}$ for the set of all paths from $(0, 0)$ to (i, j) and let $T(i, j)$ denote the weight of the maximal path from the origin to (i, j) ,

$$T(i, j) = \max_{\gamma \in \Gamma_{i,j}} \sum_{(i', j') \in \gamma} Y_{i'j'}.$$

We set $\mathcal{W}' = \{(x, y) \in \mathbb{R}^2 : y \geq 0, x + y \geq 0\}$. For $(x, y) \in \mathcal{W}'$ set

$$\tau(x, y) = \lim_{n \rightarrow \infty} \frac{1}{n} T(\lfloor xn \rfloor, \lfloor yn \rfloor).$$

In [2] it is shown that this limit exists almost surely and is deterministic (and in particular does not depend on the disorder α). It is concave, superadditive and positively 1-homogeneous. We write

$$h(t, x) = \inf\{y \geq 0 : \tau(x, y) > t\}$$

which by homogeneity can be written as $h(t, x) = tk(x/t)$. In the case of the homogeneous TASEP with constant rate r we have

$$\tau(x, y) = \frac{1}{r} (\sqrt{x+y} + \sqrt{y})^2, \quad k(v) = \begin{cases} \frac{r(1-v/r)^2}{4} & v \in [-1, 1] \\ -v & v \leq -1 \end{cases}$$

It is shown in [2] that the flux function satisfies

$$f(\rho) = \inf_{v \in \mathbb{R}} [k(v) + v\rho].$$

To analyse τ and h we will couple the process with a homogeneous rate r TASEP. Let U_{ij} be independent random variables, independent of the Y_{ij} , with law $\text{Ber}(1 - \frac{r}{\alpha(i)}) \cdot \exp(r)$, that is the product of a Bernoulli and an exponential random variable. Let $Z_{ij} = Y_{ij} + U_{ij}$ which is an exponential with rate r . Thus $\mathbb{E}Y_{ij} = \frac{1}{r} - \mathbb{E}[\frac{1}{\alpha(i)}] = \mu > 0$. Let $\gamma_\star$ be the maximal path in the disorder system and so

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \max_{\gamma \in \Gamma_{i,j}} \sum_{(i', j') \in \gamma} Y_{i'j'} &= \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \max_{\gamma \in \Gamma_{i,j}} \sum_{(i', j') \in \gamma} Z_{i'j'} - \sum_{(i', j') \in \gamma} U_{i'j'} \\ &\leq \frac{1}{r} (\sqrt{x+y} + \sqrt{y})^2 - \liminf_n \frac{1}{n} \mathbb{E} \left[\sum_{(i', j') \in \gamma_\star} U_{i'j'} \right]. \end{aligned}$$

Let $\mathcal{F}$ be the σ -algebra generated by the disorder $(\alpha(i))$ and by Y_{ij} . Since the U_{ij} are conditionally independent given $\mathcal{F}$ almost surely we have that

$$\begin{aligned} \liminf_n \frac{1}{n} \mathbb{E} \left[\sum_{(i',j') \in \gamma_\star} U_{i'j'} \right] &= \liminf_n \frac{1}{n} \mathbb{E} [\mathbb{E} \left[\sum_{(i',j') \in \gamma_\star} U_{i'j'} \mid \mathcal{F} \right]] \\ &= \liminf_n \frac{1}{n} \mathbb{E} \left[\sum_{(i',j') \in \gamma_\star} \mathbb{E}[U_{i'j'} \mid \mathcal{F}] \right] \\ &\geq \liminf_n \frac{1}{n} \mathbb{E} \left[\sum_{i=0}^{\lfloor nx \rfloor} \mathbb{E}[U_{i0} \mid \mathcal{F}] \right] = \mu|x|. \end{aligned}$$

where the second equality uses the fact that $\gamma_\star$ is $\mathcal{F}$ measurable. The inequality uses the fact that the $\mathbb{E}[U_{i'j'} \mid \mathcal{F}]$ are positive and depend only on $\alpha(i)$ and that there is at least one $(i',j') \in \gamma_\star$ for each $i' \in \{0, \dots, \lfloor nx \rfloor\}$. The final equality is the law of large numbers applied to $\mathbb{E}[U_{i0} \mid \mathcal{F}]$ which are bounded and independent and identically distributed. Thus writing

$$\tilde{\tau}(x, y) = \frac{1}{r} (\sqrt{x+y} + \sqrt{y})^2 - \mu|x|$$

we have that

$$\tau(x, y) \leq \tilde{\tau}(x, y).$$

Now checking that

$$f(\rho) = \inf_{v \in \mathbb{R}} [k(v) + v\rho] \geq r/4$$

is equivalent to

$$\begin{aligned} \inf_{v \in \mathbb{R}} [k(v) + v\rho] \geq r/4 &\iff \inf_{v \in \mathbb{R}} \frac{4}{r} k(v) \geq 1 - \frac{4}{r} v\rho \\ &\iff \inf_{x \in \mathbb{R}} \frac{4}{r} k\left(\frac{xr}{4}\right) \geq 1 - x\rho \\ &\iff \inf_{x \in \mathbb{R}} h\left(\frac{4}{r}, x\right) \geq 1 - x\rho \\ &\iff \max_x \tau(x, 1 - x\rho) \leq \frac{4}{r}, \end{aligned} \tag{0.1}$$

where in the second line we made the substitution $v = \frac{xr}{4}$. We will show that $\max_x \tilde{\tau}(x, 1 - x\rho) \leq \tilde{\tau}(0, 1) = \frac{4}{r}$ for $\rho \in [\frac{1}{2} - \frac{1}{4}\mu r, \frac{1}{2} + \frac{1}{4}\mu r]$. Note that the left and right derivatives of $\tilde{\tau}(x, 1 - x\rho)$ are respectively

$$\lim_{h \nearrow 0} \frac{\tilde{\tau}(h, 1 - h\rho) - \tilde{\tau}(0, 1)}{h} = \frac{2 - 4\rho}{r} + \mu,$$

and

$$\lim_{h \searrow 0} \frac{\tilde{\tau}(h, 1 - h\rho) - \tilde{\tau}(0, 1)}{h} = \frac{2 - 4\rho}{r} - \mu.$$

Thus the left derivative is non-negative and the right derivative is non-positive at $x = 0$ when $\rho \in [\frac{1}{2} - \frac{1}{4}\mu r, \frac{1}{2} + \frac{1}{4}\mu r]$ then $\tau(x, 1 - x\rho)$ is maximized at $x = 0$ since it is concave establishing equation (0.1).

Thus

$$f(\rho) = \frac{r}{4} \quad \text{for } \rho \in [\frac{1}{2} - \frac{1}{4}\mu r, \frac{1}{2} + \frac{1}{4}\mu r].$$

REFERENCES

- [1] C. Bahadoran and T. Bodineau, *Existence of a plateau for the flux of tasep with site disorder*, arXiv preprint arXiv:1602.06718 (2016).
- [2] T. Seppäläinen et al., *Existence of hydrodynamics for the totally asymmetric simple k -exclusion process*, The Annals of Probability **27** (1999), no. 1, 361–415.

ALLAN SLY

DEPARTMENT OF MATHEMATICS, FINE HALL, PRINCETON UNIVERSITY, PRINCETON, NJ 08540, USA.

DEPARTMENT OF STATISTICS, UC BERKELEY, BERKELEY, CA 94720, USA.

E-mail address: `asly@math.princeton.edu`